
Appendix C. Source and Reliability of Estimates

SOURCE OF DATA

The data were collected during the first eight interviews of the 1984 panel of the Survey of Income and Program Participation (SIPP). The SIPP universe is the noninstitutionalized resident population living in the United States. This population includes persons living in group quarters, such as dormitories, rooming houses, and religious group dwellings. Crew members of merchant vessels, Armed Forces personnel living in military barracks, and institutionalized persons, such as correctional facility inmates and nursing home residents, were not eligible to be in the survey. Also, U.S. citizens residing abroad were not eligible to be in the survey. Foreign visitors who work or attend school in this country and their families were eligible; all others were not eligible. With the exceptions noted above, persons who were at least 15 years of age at the time of the interview were eligible to be interviewed in the survey.

The 1984 panel SIPP sample is located in 174 areas comprising 450 counties (including one partial county) and independent cities. Within these areas, clusters of two to four living quarters were systematically selected from lists of addresses prepared for the 1970 decennial census to form the bulk of the sample. To account for living quarters built within each of the sample areas after the 1970 census, a sample was drawn of permits issued for construction of residential living quarters through March 1983. In jurisdictions that do not issue building permits, small land areas were sampled and the living quarters within were listed by field personnel and then subsampled. In addition, sample living quarters were selected from a supplemental frame that included new construction for which building permits were issued prior to January 1, 1970, but for which construction was not completed until after April 1, 1970.

The first interview of this panel was conducted during October, November, and December 1983, and January 1984. Approximately one-fourth of the sample was interviewed in each of these months. Each sample person was visited every 4 months thereafter. At each interview the reference period was the 4 months preceding the interview month.

Approximately 26,000 living quarters were originally designated for the sample. At the first interview, interviews were obtained from the occupants of about 19,900 of the 26,000 designated living quarters. Most of the remaining 6,100 living quarters were found to be

vacant, demolished, converted to nonresidential use, or otherwise ineligible for the survey. However, approximately 1,000 of the 6,100 living quarters were not interviewed because the occupants refused to be interviewed, could not be found at home, were temporarily absent, or were otherwise unavailable. Thus, occupants of about 95 percent of all eligible living quarters participated in the first interview of the survey.

For subsequent interviews, only original sample persons (those interviewed in the first interview) and persons living with them were eligible to be interviewed. Original sample persons were followed if they moved to a new address, unless the new address was more than 100 miles from a SIPP sample area. Then, telephone interviews were attempted. All first interview noninterviewed households were automatically designated as noninterviews for all subsequent interviews. When original sample persons moved to remote parts of the country, moved without leaving a forwarding address or refused to be interviewed, additional noninterviews resulted.

A person was classified as interviewed or noninterviewed for the panel based on the following definitions. Interviewed sample persons were defined to be 1) those for whom self or proxy responses were obtained for each reference month of all eight interviews or 2) those for whom self or proxy responses were obtained for the first reference month of the panel and for each subsequent reference month until they were known to have died or moved to an ineligible address (foreign living quarters, institutions, or military barracks). Noninterviewed persons were defined to be those for whom neither self nor proxy responses were obtained for one or more reference months of the eight interviews (but not because they were deceased or moved to an ineligible address). All members of a household were excluded from the weighting procedure if one or more members had no self or proxy responses for the first interview. (The processing system was unable to handle persons in this type of household.)

Approximately 52,800 persons were counted as initially interviewed. This count excludes about 1,300 interviewed persons who were members of households in which one or more members were noninterviews in the first interview. In the weighting procedure, approximately 32,400 persons were classified as interviewed. Persons who missed interviews due to the March 1985 sample cut were not classified as noninterviews but were adjusted for in the weighting procedure by a

special factor (see "Estimation"). The person nonresponse rate is estimated to be 30 percent for the panel. Some respondents did not respond to some of the questions; therefore, the overall nonresponse rate for some items, especially sensitive income and money related items, is higher than the person nonresponse rate.

ESTIMATION

Several stages of weight adjustments were involved in the estimation procedure used to derive the SIPP longitudinal person weights. Each person received a base weight equal to the inverse of his/her probability of selection. Two noninterview adjustment factors were applied. One adjusted the weights of interviewed persons in interviewed households to account for households which were eligible for the sample but could not be interviewed at the first interview. The second was applied to compensate for person noninterviews occurring in subsequent interviews. Another factor was applied to each interviewed person's weight to account for the SIPP sample areas not having the same population distribution as the strata from which they were selected.

An additional stage of adjustment to longitudinal person weights was performed to reduce the mean square error of the survey estimates. This was accomplished by bringing the sample estimates into agreement with monthly Current Population Survey (CPS) type estimates of the civilian (and some military) noninstitutional population of the United States by age, sex, race, Hispanic ethnicity, and householder/not householder status as of the specified control date. The control date for the weights was November 1, 1983. The CPS estimates were themselves brought into agreement with estimates from the 1980 decennial census which have been adjusted to reflect births, deaths, immigration, emigration, and changes in the Armed Forces since 1980.

RELIABILITY OF ESTIMATES

SIPP estimates are based on a sample; they may differ somewhat from the figures that would have been obtained if a complete census had been taken using the same questionnaire, instructions, and enumerators. There are two types of errors possible in an estimate based on a sample survey: nonsampling and sampling. We are able to provide estimates of the magnitude of SIPP sampling error, but this is not true of nonsampling error. Found in the next sections are descriptions of sources of SIPP nonsampling error, followed by a discussion of sampling error, its estimation, and its use in data analysis.

Nonsampling variability. Nonsampling errors can be attributed to many sources, e.g., inability to obtain information about all cases in the sample, definitional difficulties, differences in the interpretation of questions,

inability or unwillingness on the part of the respondents to provide correct information, inability to recall information, errors made in collection such as in recording or coding the data, errors made in processing the data, errors made in estimating values for missing data, biases resulting from the differing recall periods caused by the interviewing pattern used, and failure of all units in the universe to have some probability of being selected for the sample (undercoverage). Quality control and edit procedures were used to reduce errors made by respondents, coders and interviewers.

Undercoverage in SIPP results from missed living quarters and missed persons within sample households. It is known that undercoverage varies with age, race, and sex. Generally, undercoverage is larger for males than for females and larger for Blacks than for non-Blacks. Ratio estimation to independent age-race-sex population controls partially corrects for the bias due to survey undercoverage. However, biases exist in the estimates to the extent that persons in missed households or missed persons in interviewed households have characteristics different from those of interviewed persons in the same age-race-sex group. Further, the independent population controls used have not been adjusted for undercoverage.

The Bureau has used complex techniques to adjust the weights for nonresponse, but the success of these techniques in avoiding bias is unknown.

Comparability with other estimates. Caution should be exercised when comparing data from this report with data from other SIPP publications or with data from other surveys. The comparability problems are caused by such sources as the seasonal patterns for many characteristics, different nonsampling errors, and different concepts and procedures.

Sampling variability. Standard errors indicate the magnitude of the sampling error. They also partially measure the effect of some nonsampling errors in response and enumeration, but do not measure any systematic biases in the data. The standard errors for the most part measure the variations that occurred by chance because a sample rather than the entire population was surveyed.

USES AND COMPUTATION OF STANDARD ERRORS

Confidence intervals. The sample estimate and its standard error enable one to construct confidence intervals, ranges that would include the average result of all possible samples with a known probability. For example, if all possible samples were selected, each of these being surveyed under essentially the same conditions and using the same sample design, and if an estimate and its standard error were calculated from each sample, then:

1. Approximately 68 percent of the intervals from one standard error below the estimate to one standard error above the estimate would include the average result of all possible samples.
2. Approximately 90 percent of the intervals from 1.6 standard errors below the estimate to 1.6 standard errors above the estimate would include the average result of all possible samples.
3. Approximately 95 percent of the intervals from two standard errors below the estimate to two standard errors above the estimate would include the average result of all possible samples.

The average estimate derived from all possible samples is or is not contained in any particular computed interval. However, for a particular sample, one can say with a specified confidence that the average estimate derived from all possible samples is included in the confidence interval.

Hypothesis testing. Standard errors may also be used for hypothesis testing, a procedure for distinguishing between population characteristics using sample estimates. The most common types of hypotheses tested are 1) the population characteristics are identical versus 2) they are different. Tests may be performed at various levels of significance, where a level of significance is the probability of concluding that the characteristics are different when, in fact, they are identical.

All statements of comparison in the report have passed a hypothesis test at the 0.10 level of significance or better. This means that, for differences cited in the report, the estimated absolute difference between parameters is greater than 1.6 times the standard error of the difference.

To perform the most common test, compute the difference $X_A - X_B$, where X_A and X_B are sample estimates of the characteristics of interest. A later section explains how to derive an estimate of the standard error of the difference $X_A - X_B$. Let that standard error be s_{DIFF} . If $X_A - X_B$ is between -1.6 times s_{DIFF} and $+1.6$ times s_{DIFF} , no conclusion about the characteristics is justified at the 10 percent significance level. If, however, $X_A - X_B$ is smaller than -1.6 times s_{DIFF} or larger than $+1.6$ times s_{DIFF} , the observed difference is significant at the 10 percent level. In this event, it is commonly accepted practice to say that the characteristics are different. Of course, some times this conclusion will be wrong. When the characteristics are, in fact, the same, there is a 10 percent chance of concluding that they are different.

Note that as more tests are performed, more erroneous significant differences will occur. For example, if 100 independent hypothesis tests are performed in which there are no real differences, it is likely that about

10 erroneous differences will occur. Therefore, if a large number of tests are performed, the significance of any single test should be interpreted cautiously.

Note concerning small estimates and small differences. Summary measures are shown in the report only when the base is 200,000 or greater. Because of the large standard errors involved, there is little chance that estimates will reveal useful information when computed on a base smaller than 200,000. Also, nonsampling error in one or more of the small number of cases providing the estimate can cause large relative error in that particular estimate. Estimated numbers are shown, however, even though the relative standard errors of these numbers are larger than those for the corresponding percentages. These smaller estimates are provided primarily to permit such combinations of the categories as serve each user's needs. Therefore, care must be taken in the interpretation of small differences since even a small amount of nonsampling error can cause a borderline difference to appear significant or not, thus distorting a seemingly valid hypothesis test.

Standard error parameters and tables and their use.

Most SIPP estimates have greater standard errors than those obtained through a simple random sample because clusters of living quarters are sampled for the SIPP. To derive standard errors that would be applicable to a wide variety of estimates and could be prepared at a moderate cost, a number of approximations were required. Estimates with similar standard error behavior were grouped together and two parameters (denoted "a" and "b") were developed to approximate the standard error behavior of each group of estimates. Because the actual standard error behavior was not identical for all estimates within a group, the standard errors computed from these parameters provide an indication of the order of magnitude of the standard error for any specific estimate. These "a" and "b" parameters vary by characteristic and by demographic subgroup to which the estimate applies. Table C-1 provides base "a" and "b" parameters to be used for estimates in this report.

For those users who wish further simplification, we have also provided general standard errors in tables C-2 and C-3. Note that these standard errors must be adjusted by a factor from table C-1. The standard errors resulting from this simplified approach are less accurate. Methods for using these parameters and tables for computation of standard errors are given in the following sections.

Standard errors of estimated numbers. The approximate standard error, s_x , of an estimated number of persons shown in this report can be obtained in two ways.

It may be obtained by the use of the formula

$$s_x = fs \quad (1)$$

Table C-1. SIPP Generalized Variance Parameters for Estimates Using Panel Weights

(1984 Longitudinal Panel File)

Persons	a	b	f factor
TOTAL OR WHITE			
15+ program participation and benefits (3):			
Both sexes	-0.0001241	22,392	.90
Male	-0.0002593	22,392	.90
Female	-0.0002380	22,392	.90
15+ income and labor force (4):			
Both sexes	-0.0000424	7,634	.52
Male	-0.0000884	7,634	.52
Female	-0.0000811	7,634	.52
All others ¹ (5):			
Both Sexes	-0.0001196	27,763	1.00
Male	-0.0002462	27,763	1.00
Female	-0.0002327	27,763	1.00
BLACK			
Poverty (1):			
Both sexes	-0.000687	19,100	.83
Male	-0.001467	19,100	.83
Female	-0.001293	19,100	.83
All others (2):			
Both sexes	-0.0003696	10,271	.61
Male	-0.0007889	10,271	.61
Female	-0.0006953	10,271	.61

¹These parameters are to be used for all tabulations not specifically covered by any other category in this table, e.g., for retirement and pension tabulations, for O+ benefits, O+ income, and O+ labor force tabulations.

Note: For cross-tabulations, apply the parameters of the category showing the smaller number in parentheses.

where f is the appropriate "f" factor from table C-1, and s is the standard error of the estimate obtained by interpolation from table C-2. Alternatively, s_x may be approximated by the formula

$$s_x = \sqrt{ax^2 + bx} \quad (2)$$

Here x is the estimated number and "a" and "b" are the parameters associated with the particular type of characteristic. Use of formula (2) will provide more accurate results than the use of formula (1).

Illustration. Suppose the SIPP estimate of the total number of persons with low income for 1 to 6 months during the 32 months of the 1984 panel is 34,894,000. The appropriate "a" and "b" parameters to use in calculating a standard error for the estimate are obtained from table C-1. They are $a = -0.0001196$, $b = 27,763$, respectively. Using formula (2), the approximate standard error is

$$\sqrt{(-0.0001196)(34,894,000)^2 + (27,763)(34,894,000)} = 907,000$$

The 90-percent confidence interval as shown by the data is from 33,443,000 to 36,345,000. Therefore, a conclusion that the average estimate derived from all

possible samples lies within a range computed in this way would be correct for roughly 90 percent of all samples.

Using formula (1), the appropriate "f" factor ($f = 1.00$) from table C-1, and the standard error of the estimate from table C-2, the approximate standard error is

$$s_x = 1.00(899,000) = 899,000$$

The 90-percent confidence interval as shown by the data is from 33,456,000 to 36,332,000.

Standard errors of estimated percentages. This section refers to the percentages of a group of persons possessing a particular attribute such as the percentage of persons participating in a major assistance program for one or more months. The reliability of an estimated percentage, computed using sample data for both numerator and denominator, depends upon both the size of the percentage and the size of the total upon which the percentage is based. Estimated percentages are relatively more reliable than the corresponding estimates of the numerators of the percentages, particularly if the percentages are over 50 percent. For example, the percent of employed persons is more reliable than the estimated number of employed persons. When the numerator and denominator of the percentage have different parameters, use the parameter (and appropriate factor) of the numerator. If proportions are presented instead of percentages, note that the standard error of a proportion is equal to the standard error of the corresponding percentage divided by 100.

For the percentage of persons, the approximate standard error, $s_{(x,p)}$, of the estimated percentage, p , can be obtained by the formula

$$s_{(x,p)} = fs \quad (3)$$

where f is the appropriate "f" factor from table C-1, and s is the standard error of the estimate from table C-3.

Table C-2. Standard Errors of Estimated Numbers of Persons

(Numbers in thousands)

Size of estimate	Standard error	Size of estimate	Standard error
200	74	50,000	1,044
300	91	80,000	1,206
600	129	100,000	1,257
1,000	166	130,000	1,260
2,000	235	135,000	1,252
5,000	369	150,000	1,214
8,000	463	160,000	1,175
11,000	539	180,000	1,059
13,000	584	200,000	877
15,000	624	210,000	746
17,000	661	220,000	565
22,000	744		
26,000	801		
30,000	852		

Table C-3. Standard Errors of Estimated Percentages of Persons

Base of estimated percentage (thousands)	Estimate percentage					
	1 or 99	2 or 98	5 or 95	10 or 90	25 or 75	50
200	3.7	5.2	8.1	11.2	16.1	15.8
300	3.0	4.3	6.6	9.1	13.2	12.9
600	2.1	3.0	4.7	6.5	9.3	9.1
1,000	1.7	2.3	3.6	5.0	7.2	7.1
2,000	1.2	1.6	2.6	3.5	5.1	5.0
5,000	0.7	1.0	1.6	2.2	3.2	3.2
8,000	0.6	0.8	1.3	1.8	2.6	2.5
11,000	0.5	0.7	1.1	1.5	2.2	2.1
13,000	0.5	0.6	1.0	1.4	2.0	2.0
17,000	0.4	0.6	0.9	1.2	1.7	1.7
22,000	0.4	0.5	0.8	1.1	1.5	1.5
26,000	0.3	0.5	0.7	1.0	1.4	1.4
30,000	0.3	0.4	0.7	0.9	1.3	1.3
50,000	0.2	0.3	0.5	0.7	1.0	1.0
80,000	0.2	0.3	0.4	0.6	0.8	0.8
100,000	0.2	0.2	0.4	0.5	0.7	0.7
130,000	0.1	0.2	0.3	0.4	0.6	0.6
220,000	0.1	0.2	0.2	0.3	0.5	0.5

Alternatively, it may be approximated by the formula

$$s_{(x,p)} = \sqrt{\frac{b}{x} p(100-p)} \quad (4)$$

Here x is the base of the percentage, p is the percentage ($0 \leq p \leq 100$), and b is the "b" parameter associated with the characteristic in the numerator. Use of this formula will give more accurate results than use of formula (3).

Illustration. Suppose that we have a SIPP estimate of 6,150,000 males aged 20 to 64 who were without a full-time job for all 32 months. Of these, 18.7 percent participated in a major assistance program for all 32 months. Using formula (4) and the "b" parameter of 22,392 (from table 1), the approximate standard error is

$$\sqrt{\frac{(22,392)}{(6,150,000)}(18.7)(100-18.7)} = 2.4 \text{ percent}$$

Consequently, the 90-percent confidence interval as shown by these data is from 14.9 to 22.5 percent.

Using formula (3), the appropriate "f" factor ($f=.90$) from table C-1, and the appropriate s from table C-3, the approximate standard error is

$$s_x = (.90)(2.6) = 2.3 \text{ percent}$$

The 90-percent confidence interval shown by these data is from 15.0 to 22.4 percent.

Standard error of a difference. The standard error of a difference between two sample estimates, x and y , is equal to

$$s_{(x-y)} = \sqrt{s_x^2 + s_y^2} \quad (5)$$

where s_x and s_y are the standard errors of the estimates x and y . The estimates can be numbers, averages, percents, ratios, etc. The above formula assumes that the correlation coefficient, r , between the characteristics estimated by x and y is zero. If r is really positive (negative), then this assumption will result in a tendency towards overestimates (underestimates) of the true standard error.

Illustration. Suppose that we are interested in the difference in the proportions of males and females aged 20 to 64 without a full-time job who participated in a major assistance program for all 32 months of the 1984 panel. An estimated 6.2 million men and 26.1 million women were without a full-time job for all 32 months. Of these, 18.7 percent and 13.6 percent, respectively, participated in a major assistance program for all the months. Using parameters and factors from table C-1, the standard errors of these numbers are approximately 2.4 percent and 1.0 percent for males and females, respectively.

Now, the standard error of the difference is computed using the above two standard errors. The standard error of the difference is computed by formula (5):

$$\sqrt{(2.4)^2 + (1.0)^2} = 2.6 \text{ percent}$$

Suppose that it is desired to test at the 10 percent significance level whether the proportion of males differs significantly from the proportion of females. To perform the test, compare the difference of 5.1 percent to the product 1.6×2.6 percent = 4.1 percent. Since the difference is larger than 1.6 times the standard error of the difference, the data show that the male estimate is significantly different from the female estimate at the 10 percent level.